

# Yet another proof of Hawking and Ellis's Lemma 8.5.5

S. Krasnikov\*

## Abstract

The fact that the null generators of a future Cauchy horizon are past complete was first proved by Hawking and Ellis [1]. Then Budzyński, Kondracki, and Królak outlined a proof free from the error found in the original one [2]. Finally, Minguzzi has recently published his version of the proof [3] patching a previously unnoticed hole in the preceding two. I am not aware of any flaws in that last proof, but it is quite difficult. In this note I present a simpler one.

## 1 Introduction

Let  $H^+$  denote a future Cauchy horizon. A lemma by Hawking and Ellis says

**Lemma 8.5.5 of [1].** If  $H^+(\mathcal{Q})$  is compact for a partial Cauchy surface  $\mathcal{Q}$ , then the null geodesic generating segments of  $H^+(\mathcal{Q})$  are geodesically complete in the past direction.

The lemma itself has never been doubted (to my knowledge), but the proof offered in [1] was found to be flawed, see [3] and references therein. To improve this situation — which is important, because the lemma is a popular tool in mathematical relativity — Minguzzi has recently published a new, more accurate, proof of the lemma (or, to be precise, of some strengthening of it).

In this comment I present yet another proof of the same fact. The reason for doing this is that my version is much simpler (partly because its major part is replaced by a reference to a lemma proved elsewhere).

---

\*Email: S.V.Krasnikov@mail.ru

## 2 The proposition and its proof

In a spacetime  $M$  consider a past inextendible null curve  $\gamma$  totally imprisoned in a compact set  $\mathcal{K}$ . Pick a smooth unit timelike future directed vector field  $\boldsymbol{\tau}$  on  $M$  and define (uniquely up to an additive constant) the “arc length parameter”  $l$  on  $\gamma$  by the requirement

$$g(\partial_l, \boldsymbol{\tau}) = -1. \quad (1)$$

In addition to  $l$  define on  $\gamma$  an *affine* parameter  $s$  so that  $\partial_s$  is future directed and  $s = 0$  at  $l = 0$ . Now  $\gamma$  is characterized by the (evidently negative) function

$$h \equiv g(\partial_s, \boldsymbol{\tau}),$$

which relates  $l$  to  $s$ :

$$h = -\frac{dl}{ds}, \quad s(l) = \int_l^0 \frac{d\check{l}}{h(\check{l})}. \quad (2)$$

As is proven in [4]

$$h'/h \text{ is bounded on } \gamma. \quad (3)$$

Further, a few minor changes — a past inextendible  $\gamma(l)$  with  $l \in (-\infty, 0]$  instead of the future inextendible  $\gamma(l)$  with  $l \in [0, \infty)$  and an arbitrary compact  $\mathcal{K}$  instead of some specific  $\mathcal{L}$  — leave [4, Lemma 8] valid while bringing it to the following form.

**Lemma 8 of [4].** Assume  $h(l)$  is such that there exists a smooth function  $f(l)$  defined at non-positive  $l$  and obeying for some non-negative constants  $c_1, \underline{f}, \bar{f}$  the inequalities

$$\underline{f} \leq f \leq \bar{f}, \quad |f'/f| < \infty$$

and

$$h'/h < -f'/f - c_1 f, \quad \forall l \leq 0. \quad (4)$$

Then there is a timelike past inextendible curve  $\gamma_{\neq 0}$  which is obtained by moving each point of  $\gamma$  to the past along the integral curves of  $\boldsymbol{\tau}$  and which is totally imprisoned in a compact set  $\mathcal{O}$ .

**Proposition.** If  $\mathcal{K}$  is a subset of the boundary of a globally hyperbolic past set  $M^{\text{in}}$ , then  $\gamma$  is past complete.

*Proof.* Suppose the lemma is false and  $\gamma$  is past incomplete. This would mean that the affine parameter  $s$  is bounded from below and, correspondingly, the integral (2) converges at  $l \rightarrow -\infty$ . Which allows one to define the following smooth positive function on  $(-\infty, 0]$

$$f(l) \equiv \frac{1}{h} \left[ -\int_l^0 \frac{d\check{l}}{h(\check{l})} + 2 \int_{-\infty}^0 \frac{d\check{l}}{h(\check{l})} \right]^{-1}. \quad (5)$$

$f$  so defined satisfies the equation

$$f'/f + h'/h = -f \quad (6)$$

and consequently, condition (4) holds.

As  $h$  is negative, the boundedness of the integral (2) provides a simple estimate

$$\infty > -\int_{-\infty}^0 \frac{d\check{l}}{h(\check{l})} > -\frac{1/h(l)}{(|h'/h|)_{\max}}, \quad \forall l \in (-\infty, 0],$$

which implies, due to (3), that  $1/h$  is bounded. It follows then from (5) that  $f$  is bounded too. Finally, the just proven boundedness of  $f$  combined with (6) and (3) implies the boundedness of  $f'/f$ . Thus all the conditions of Lemma 2 are fulfilled and the corresponding variation transforms  $\gamma$  into a past inextendible timelike curve  $\gamma_{\varkappa_0}$ . The latter being timelike lies entirely in the closed (due to the globally hyperbolicity of  $M^{\text{in}}$ , to which  $\gamma_{\varkappa_0}(0)$  belongs) set  $J^-(\gamma_{\varkappa_0}(0)) \subset M^{\text{in}}$  (the inclusion follows from the fact that  $M^{\text{in}}$  is a past set). Thus,  $\gamma_{\varkappa_0}$  is totally imprisoned in the compact subset  $\mathcal{O} \cap J^-(\gamma_{\varkappa_0}(0))$  of the globally hyperbolic spacetime  $M^{\text{in}}$ , which is forbidden by [1, proposition 6.4.7].

□

## References

- [1] S. W. Hawking and G. F. R. Ellis *The Large scale structure of spacetime* (Cambridge, Cambridge University Press, 1973).
- [2] R. Budzyński, W. Kondracki, A. Krolak *New properties of Cauchy and event horizons*, arXiv:gr-qc/0011033.

- [3] E. Minguzzi, J. Math. Phys. **55**, 082503 (2014).
- [4] S. Krasnikov, Physical Review D **90**, 024067 (2014).